



Hydrogeochemical Parameters for Assessment of Groundwater Quality in Marbache Ramhormoz, Khouzestan, Iran

Payvad Papan¹, Nooshin Shahinzadeh², Ali Gholami³

¹KWPA- Khouzestan Water and Power Authority CO.

²Ph.D Student, Department of Soil Science, Science and Research Branch, Islamic Azad University, Khouzestan, Iran.

³Assistant Professor of Soil Science, Science and Research Branch, Islamic Azad University, Khouzestan, Iran

Abstract: In the management of water resources, quality of water is just as important as its quantity. In order to know the quality and/or suitability of groundwater for irrigation in ramhormoz, 13 water samples dry season were collected and analyzed for various parameters. Based on the analytical results, chemical indices like percent sodium, sodium adsorption ratio, electrical conductivity, chloride, pH, sodium, calcium, magnesium, bicarbonate, total dissolved solids, sulfate, were calculated. The parameters were used to assess the suitability of groundwater for irrigation purpose by comparing with FAO. The sample analysis results that the groundwater is entirely fit for agriculture.

Keywords: Hydrochemical parameters, groundwater quality, SAR, Marbache Ramhormoz,

1. Introduction

In recent times, there are been a tremendous increase in demand for fresh water due to population growth and intense agricultural activities. Groundwater is principal source of irrigation water in our country and indispensable source of our life. The quality of the groundwater varies from place to place



۲۹ و۳۰ اردیبهشت ماه ۱۳۹۴ - مرکز همایش های بین المللی شهید بهشتی 19 - 20 May, 2015 - Shahid Beheshti Conf. Center

with the depth of water table. The classification, modeling and interpretations of monitoring data are the most important steps in the assessment of water quality causes. It is a very difficult and laborious task to regularly monitor all the parameters even if adequate manpower and laboratory facilities are available. For this reason, in recent years an easier and simpler approach based on statistical correlation, has been developed using mathematical relationship for comparison of chemical parameters (lyer, C.S et al., 2003; Sarkar Mitali et al., 2006). The correlation coefficient is a helpful tool for the promotion of research in water quality.

2. Materials and Methods

2.1 Study area

Ramhormoz located in the Khouzestan plain at the latitude of 31 10' to 31° 20'north and longitude of 49° 15' to 49° 45' west covering an area of approximately 500 sq. km.

2.2 Methods

A total of eightheen groundwater samples were collected in 2.5 liter polyethylene bottles during dry season in July 2013. During the present study, the samples were analyzed for various chemical parameters as described by the American Public Health Association (APHA,1995). These parameters include pH, electrical conductivity (EC), total dissolved solids (TDS), and important cations such as calcium, magnesium, sodium and potassium as well as anions such as bicarbonates, chlorides, sulfates.



۲۹ و ۳۰ اردیبهشت ماه ۱۳۹۴ - مرکز همایش های بین المللی شهید بهشتی 19 - 20 May, 2015 - Shahid Beheshti Conf. Center

The Ph and EC were measured by means of pH meter (Metrohm pH 691) and conductivity meter (Session Con.378). The TDS of the samples were also computed from conductivity meter. Sodium and potassium were determined by flame photometer (Flame photometer 416). Total hardness as CaCO₃, Calcium (Ca²⁺), magnesium (Mg²⁺), bicarbonate (HCO₃⁻) and chloride (Cl⁻) were analyzed by titrimetry method. Sulfate (SO₄²⁻) was analyzed by spectrophotometer. The statistical analysis has been performed to evaluate correlation coefficients between different pairs of water quality parameters. The linear regression approach was also carried out to quantify the relationship between several independent or predictor variables and dependent variable.

2.3 Statistical analysis

2.3.1 Pearson correlation

Pearson correlation coefficient is commonly used to measure and establish the strength of a linear relationship between two variables or two sets of data. It is a simplified statistical tool to show the degree of dependency of one variable to the other (Belkhiri et al., 2010).

2.3.2 Multiple linear regressions

In this study, we have applied the multiple linear regression approach to develop a relationship between several independent variables and dependent variable.

3. Results and Discussion

The minimum and maximum concentration i.e. the range of the different chemical parameters of water quality such as pH, electrical conductivity (EC), total dissolved solid (TDS), sodium (Na), potassium (K), sulphate ($SO_4^{2^-}$), calcium (Ca), magnesium (Mg), chloride (Cl) in the study area given in Table 1 along with Mean, Standard Deviation, Variance and Standard Error for dry season.



بایشگاه مهند م

۲۹ و۳۰ اردیبهشت ماه ۱۳۹۴ - مرکز همایش های بینالمللی شهید بهشتی 19 - 20 May, 2015 - Shahid Beheshti Conf. Center

No.	parameter	Min. (mg/l)	Max.(mg/l)	Mean	Std.error	Std. Deviation	Variance
1	EC	1550	9250	4204.84	659.40	2377.52	565263
2	рН	6.3	7.90	7.18	0.11921	0.4298	0.185
3	TDS	1163	7404	3047.46	490.56	1768.76	3128516
4	Ca ²⁺	4.70	34.63	18.35	2.15	7.78	60.609
5	Mg ²⁺	3.37	25.27	10.49	1.87	6.77	45.944
6	Na⁺	4.53	57	16.95	4.25	15.33	235.093
7	K+	0.07	1.30	0.3238	0.1239	0.4468	0.2
8	HCO₃ ⁻	1.70	6.10	3.05	0.35154	1.26	1.60
9	Cl-	3.35	58.19	17.54	4.19	15.11	228.447

Table 1: Basic statistics of groundwater in July 2013

	Eng. Vater				- مایش های بینالمللی شهید	می نمایشگاه ک و نمایشگاه م بهشت ماه ۱۳۹۴- مرکز ه hahid Beheshti Cont	۲۹ و۳۰ ارد.	
10	SO4 ²⁻	6.91	58.73	25.27	4.10	14.80	219.196	

Table 1 indicates that the pH values of groundwater in the study area range from 6.8-7.5 in dry seasons. The pH values of all samples are within the desirable limit (6.5-8.5). TDS is an important parameter for assessing groundwater quality. All samples in dry season cross the permissible limit for TDS (500 mg/l-2000mg/l). The EC values of all samples are within the desirable limit (0.7 ds/m- 3ds/m). Most of the samples fall within the desirable limit for Cl (250 mg/l).

The correlation (r) measures the degree of association that exists between two variables one taken as dependent variable. It is evident from Table 2 that during dry season, highly positive correlation coefficient characterized by r = 0.8 to 1.0 is observed between EC- Ca, EC-Na, EC- CL and TDS-Na, TDS-Ca, TDS- CL, TDS- EC, Na- Cl, Ca- SO4, Mg- SO4. As indicated in Table 2, pH has little association with most of the water quality parameters in dry season.

parameter	EC	рН	TDS	Ca ²⁺	Mg ²⁺	Na⁺	K+	HCO₃ ⁻	Cl-	SO4 ²⁻
EC	1									
рН	0.236	1								
TDS	0.977	0.109	1							
Ca ²⁺	0.875	0.1	0.842	1						
Mg ²⁺	0.638	0.076	0.627	0.787	1					
Na⁺	0.912	0.312	0.871	0.659	0.308	1				
K+	0.650	-0.073	0.727	0.513	0.673	0.433	1			
HCO3 ⁻	0.788	-0.331	0.494	0.371	0.348	0.187	0.379	1		
CI-	0.879	0.343	0.838	0.608	0.226	0.994	0.384	0.155	1	

wate	ig.							المللي شهيد بهش	نو میایش های بیز 2015 - Shał	ماه ۱۳۹۴-مرک	۳۰ اردیبهشت	۲۹و
SO4 ²⁻	0.788	0.87	0.756	0.932	0.950	0.504	0.631	0.317	0.429	1		

Regression analysis was conducted to investigate the relationships between TDS and other water properties, which shows significant correlation with TDS (r = 0.8 to 1) using SPSS. The [Cl], [EC], [Na], [Ca] were considered as independent variables and TDS as a dependent variable. An analysis of residuals was developed and R² values were studied. Among all candidate equations, the equation where this ratio was closer to 1 was selected. The descriptors and the regression coefficient of this model are presented in Table3. The positive sign of the beta coefficient pertaining to these variables indicates that there is a positive relationship between TDS and elements of groundwater properties[Na], [Ca], [Cl], [EC]. The selected regression equation is given by:

TDS= 49.001 + 1.08 EC-52.72 Ca-35.36 Na + 0.54 Cl

Regression	Coefficients
βΟ	49.001
EC (β1)	1.08
Cl (β2)	0.54
Na (β3)	-35.36
Ca (β4)	-52.72

Table 3: Summary of multiple linear regressions predicting TDS





۲۹ و۳۰ اردیبهشت ماه ۱۳۹۴ - مرکز همایش های بینالمللی شهید بهشتی 19 - 20 May, 2015 - Shahid Beheshti Conf. Center

Correlation	Coefficient
Multiple R	1.000
R square	0.963
Adjusted R Square	0.944

The actual and predicted value for TDS for dry season was plotted in a graph (Fig 1)



Figure 1: Actual Vs predicted values for TDS (dry season)

4. Conclusion

Multivariate statistical method used in this study (Pearson correlation coefficients and multiple linear regressions) help to find statistically important factors in data variability and thus improve conclusions in environmental impact studies. Pearson correlation matrix was applied to all the collected water samples for identifying the possible statistical relationship between different pairs of ground water quality parameters. A highly strong correlation was observed between Ca²⁺ and HCO₃⁻ in dry season, which gives





us an idea about the total hardness of water. A multiple linear regression was used to establish relationship between TDS and other chemical water properties. The positive sign of the regression coefficient indicates that there is a positive relationship between TDS and elements of groundwater properties: [Na], [Ca], [Cl], [EC]. It can be concluded that the total dissolved solids and electrical with most of the elements in the groundwater.

5. Acknowledgements

Thanks to the Research and Standards Office for Irrigation and Drainage Networks of KWPA for Financial supports and all assistance.

6. References

1. APHA (1995), Standard methods for the examination of water waste water, 19th edn. American Public Health Association, Washington, DC.

2. Belkhiri, L., Boudoukha, A., Mouni, L., (2010), A multivariate Statistical Analysis of Groundwater Chemistry Data, International Journal of Environmental Research, 5(2), 537-544.



و نمایشگاه مهند آب كنفر

۲۹ و۳۰ ارديبهشت ماه ۱۳۹۴ - مرکز همايشرهای بيزالمللی شهيد بهشتی 19 - 20 May, 2015 - Shahid Beheshti Conf. Center